

The Physics of Information Technology

Neil Gershenfeld

revised draft: February 9, 2022

© Cambridge University Press

Contents

<i>Preface</i>	<i>page</i> ix x
1 Introduction	1
2 Interactions, Units, and Magnitudes	3
2.1 Units	3
2.2 Particles and Forces	7
2.3 Orders of Magnitude	10
2.4 Selected References	12
2.5 Problems	12
3 Noise in Physical Systems	14
3.1 Random Variables	14
3.1.1 Expectation Values	14
3.1.2 Spectral Theorems	15
3.2 Probability Distributions	18
3.2.1 Binomial	18
3.2.2 Poisson	18
3.2.3 Gaussian	19
3.2.4 Central Limit Theorem	21
3.3 Noise Mechanisms	22
3.3.1 Shot Noise	22
3.3.2 Johnson Noise	23
3.3.3 $1/f$ Noise and Switching Noise	24
3.3.4 Amplifier Noise	25
3.4 Thermodynamics and Noise	27
3.4.1 Thermodynamics and Statistical Mechanics	27
3.4.2 Equipartition Theorem	30
3.4.3 Fluctuation–Dissipation Theorem	31
3.5 Selected References	34
3.6 Problems	35
4 Information in Physical Systems	36
4.1 Information	36

4.2	Channel Capacity	40
4.3	The Gaussian Channel	42
4.4	Fisher Information	45
4.5	Information and Thermodynamics	48
4.6	Selected References	49
4.7	Problems	49
5	Logic and Computation	51
5.1	Logic	53
5.2	Computation	53
5.3	Problems	53
6	Electromagnetic Fields and Waves	54
6.1	Vector Calculus	54
6.1.1	Vectors	54
6.1.2	Differential Operators	56
6.1.3	Integral Relationships	58
6.2	Statics	59
6.2.1	Electrostatics	59
6.2.2	Magnetostatics	63
6.2.3	Multipoles	66
6.3	Dynamics	69
6.3.1	Maxwell's Equations	69
6.3.2	Boundary Conditions	71
6.3.3	Electromagnetic Units	72
6.4	Radiation and Energy	74
6.4.1	Waves	74
6.4.2	Electromagnetic Energy	76
6.5	Selected References	77
6.6	Problems	77
7	Circuits, Transmission Lines, and Waveguides	80
7.1	Circuits	80
7.1.1	Current and Voltage	80
7.1.2	Kirchhoff's Laws	81
7.1.3	Resistance	81
7.1.4	Power	83
7.1.5	Capacitance	83
7.1.6	Inductance	84
7.2	Wires and Transmission Lines	85
7.2.1	Skin Depth	85
7.2.2	Transmission Lines	87
7.2.3	Wave Solutions	89
7.2.4	Reflections and Terminations	91
7.3	Waveguides	94
7.3.1	Governing Equations	94

7.3.2 Rectangular Waveguides	96
7.3.3 Circular Waveguides	97
7.3.4 Dielectric Waveguides and Fiber Optics	97
7.4 Selected References	99
7.5 Problems	100
8 Antennas	102
8.1 Time-Dependent Potentials	102
8.2 Dipole Radiation	106
8.2.1 Infinitesimal Length	106
8.2.2 Finite Length	108
8.3 Duality and Reciprocity	110
8.4 Antenna Types	113
8.5 Selected References	115
8.6 Problems	115
9 Optics	116
9.1 Reflection and Refraction	116
9.2 Geometrical Optics	121
9.2.1 Ray Matrices	123
9.2.2 Optical Transforms	124
9.3 Beyond Geometrical Optics	126
9.3.1 Gaussian Optics	126
9.3.2 Nonlinear Optics	127
9.4 Metamaterials	129
9.4.1 Negative Indices	129
9.4.2 Anomalous Dispersion	129
9.4.3 Perfect Lenses	131
9.4.4 Cloaking	131
9.5 Selected References	132
9.6 Problems	132
10 Lensless Imaging and Inverse Problems	134
10.1 Matched Filters and Synthetic Lenses	134
10.2 Coherent Imaging	137
10.3 Computed Tomography	140
10.4 Magnetic Resonance Imaging	141
10.5 Inverse Problems	143
10.6 Computational Imaging	145
10.6.1 Confocal	145
10.6.2 Near-Field	146
10.6.3 Ultrafast	146
10.6.4 Single-Pixel	146
10.6.5 Light Fields	146
10.7 Selected References	146
10.8 Problems	146

11 Semiconductor Materials and Devices	148
11.1 Quantum Statistical Mechanics	148
11.2 Electronic Structure	150
11.3 Junctions, Diodes, and Transistors	158
11.4 Logic	163
11.5 Limits	168
11.6 Selected References	171
11.7 Problems	171
12 Optical Materials and Devices	173
12.1 Generation	173
12.1.1 Incandescence	173
12.1.2 Luminescence	175
12.2 Detection	179
12.3 Modulation	183
12.3.1 Polarization	184
12.3.2 Liquid Crystals	186
12.3.3 AOM, EOM	189
12.3.4 Smoke and Mirrors	191
12.4 Selected References	194
12.5 Problems	195
13 Magnetic Materials and Devices	196
13.1 Magnetism	196
13.1.1 Diamagnetism	198
13.1.2 Paramagnetism	199
13.1.3 Ferro-, antiferro-, and ferri-magnetism	201
13.2 Magnetic Recording	206
13.2.1 Magnetic Media	206
13.2.2 Magnetic Recording	207
13.2.3 Recording Systems	209
13.3 Hall Effect	210
13.3.1 Classical	210
13.3.2 Quantum	211
13.4 Spintronics	211
13.5 Selected References	213
13.6 Problems	213
14 Transducers	214
14.1 Many-Body Effects	214
14.1.1 Superconductivity	215
14.1.2 Junctions and Interferometers	218
14.2 Non-Equilibrium Thermodynamics	220
14.2.1 Thermoelectricity	222
14.2.2 Piezoelectricity	225
14.3 Relativity	226

14.3.1	Clocks	226
14.3.2	Time	229
14.3.3	Position	232
14.4	Selected References	234
14.5	Problems	235
15	Measurement and Coding	236
15.1	Instrumentation	236
15.1.1	Amplifiers	236
15.1.2	Operational Amplifiers	236
15.1.3	Grounding, Shielding, and Leads	240
15.1.4	Bridges	242
15.1.5	Network Analyzers	243
15.2	Modulation and Detection	243
15.2.1	Synchronous Detection	244
15.2.2	Phase Detection and Encoding	246
15.2.3	Spread Spectrum	249
15.2.4	Digitization	253
15.3	Coding	254
15.3.1	Compression	255
15.3.2	Error Correction	257
15.3.3	Channel Coding	259
15.3.4	Cryptography	260
15.4	Sensing	261
15.4.1	Analog Logic	261
15.4.2	Compressed Sensing	262
15.5	Selected References	263
15.6	Problems	264
16	Quantum Computing and Communications	266
16.1	Quantum Mechanics	267
16.1.1	States and Operators	267
16.1.2	Angular Momentum	273
16.1.3	Density Matrices	277
16.2	Information	280
16.3	Communications	283
16.3.1	Cryptography	283
16.3.2	Circuits	284
16.3.3	Teleportation	285
16.3.4	Error Correction	287
16.4	Computation	289
16.4.1	Searching	290
16.4.2	Transforms and Factoring	292
16.4.3	Simulation	295
16.4.4	Optimization	296
16.4.5	Experimental Implementation	296

16.5 Selected References	298
16.6 Problems	299
<i>Appendix 1 Problem Solutions</i>	301
A1.1 Introduction	301
A1.2 Interactions, Units, and Magnitudes	301
A1.3 Noise in Physical Systems	307
A1.4 Information in Physical Systems	313
A1.5 Electromagnetic Fields and Waves	318
A1.6 Circuits, Transmission Lines, and Wave Guides	326
A1.7 Antennas	330
A1.8 Optics	333
A1.9 Lensless Imaging and Inverse Problems	340
A1.10 Semiconductor Materials and Devices	346
A1.11 Optical Materials and Devices	351
A1.12 Magnetic Materials and Devices	356
A1.13 Transducers	360
A1.14 Measurement and Coding	364
A1.15 Quantum Computing and Communication	374
 <i>Bibliography</i>	 379
<i>Index</i>	396

2 Interactions, Units, and Magnitudes

Modern information technology operates over a spectacular range of scales; bits from a memory cell with a size of 10^{-8} meters might be sent 10^7 meters to a geosynchronous satellite. It is important to be comfortable with the orders of magnitudes and associated interaction mechanisms that are useful in practice. Our first task will be to review the definitions of important units, then survey the types of forces, and finally look at typical numbers in various regimes.

2.1 UNITS

Many powers of ten have been named because it is much easier to say something like “a femtosecond optical pulse” than “a 0.000 000 000 000 001 second optical pulse” when referring to typical phenomena at that scale (a cycle of light takes on the order of a femtosecond). The dizzying growth of our ability to work with large and small systems pushes the bounds of this nomenclature; data from exabyte storage systems is read out into femtofarad memory cells. It is well worth memorizing the prefixes in Table 2.1.

Table 2.1. *Orders of magnitude.*

Magnitude	Prefix	Symbol	Magnitude	Prefix	Symbol
10^{-24}	yocto	y	10^{24}	yotta	Y
10^{-21}	zepto	z	10^{21}	zetta	Z
10^{-18}	atto	a	10^{18}	exa	E
10^{-15}	femto	f	10^{15}	peta	P
10^{-12}	pico	p	10^{12}	tera	T
10^{-9}	nano	n	10^9	giga	G
10^{-6}	micro	μ	10^6	mega	M
10^{-3}	milli	m	10^3	kilo	k
10^{-2}	centi	c	10^2	hecto	h
10^{-1}	deci	d	10^1	deka	da

Physical quantities must of course be measured in a system of units; there are many alternatives that are matched to different regimes and applications. Because of their inter-relationships it is necessary only to define a small number of fundamental quantities to

be able to derive all of the other ones. The choice of which fundamental definitions to use changes over time to reflect technological progress; once atomic clocks made it possible to measure time with great *precision* (small variance) and *accuracy* (small bias), it became more reliable to define the meter in terms of time and the speed of light rather than a reference bar kept at the Bureau International des Poids et Mesures (BIPM, <http://www.bipm.org>) in Sevres, France [Giacomo, 1984]. More recently, a great source of frustration in the metrology community [Girard, 1994] was removed when the definition of the kilogram based on a platinum–iridium cylinder held at BIPM was replaced by one based on fundamental physical constants [Stock *et al.*, 2019].

The most common set of base defined quantities in use is the *Système International d’Unités (SI)* [BIPM, 2019]. It is based on seven physical constants that were originally measured and are now taken as fixed (there is no uncertainty in these values):

- $\Delta\nu_{\text{Cs}} = 9\,192\,631\,770$ Hz: the unperturbed ground state hyperfine transition frequency of the cesium 133 atom
- $c = 299\,792\,458$ m/s: the speed of light in vacuum
- $h = 6.626\,070\,15 \times 10^{-34}$ J·s: the Planck constant
- $e = 1.602\,176\,634 \times 10^{-19}$ C: the elementary charge
- $k = 1.380\,649 \times 10^{-23}$ J/K: the Boltzmann constant
- $N_{\text{A}} = 6.022\,140\,76 \times 10^{23}$ mol⁻¹: the Avogadro constant
- $K_{\text{cd}} = 683$ lm/W: the luminous efficacy of monochromatic radiation of frequency 540×10^{12} Hz

These constants then define seven base units (which were previously themselves taken as fundamental):

time: *second* (s)

It is defined by taking the fixed numerical value of the cesium frequency $\Delta\nu_{\text{Cs}}$, the unperturbed ground-state hyperfine transition frequency of the cesium 133 atom, to be 9 192 631 770 when expressed in the unit Hz, which is equal to s⁻¹.

length: *meter* (m)

It is defined by taking the fixed numerical value of the speed of light in vacuum c to be 299 792 458 when expressed in the unit m·s⁻¹, where the second is defined in terms of $\Delta\nu_{\text{Cs}}$.

mass: *kilogram* (kg)

It is defined by taking the fixed numerical value of the Planck constant h to be $6.626\,070\,15 \times 10^{-34}$ when expressed in the unit J·s, which is equal to kg·m²·s⁻¹, where the metre and the second are defined in terms of c and $\Delta\nu_{\text{Cs}}$.

electric current: *ampere* (A)

It is defined by taking the fixed numerical value of the elementary charge e to be $1.602\,176\,634 \times 10^{-19}$ when expressed in the unit C, which is equal to A·s, where the second is defined in terms of $\Delta\nu_{\text{Cs}}$.

temperature: *kelvin* (K)

It is defined by taking the fixed numerical value of the Boltzmann constant k to be $1.380\,649 \times 10^{-23}$ when expressed in the unit J·K⁻¹, which is equal to kg·m²·s⁻²·K⁻¹, where the kilogram, meter and second are defined in terms of h , c and $\Delta\nu_{\text{Cs}}$.

amount of substance: *mole* (mol)

One mole contains exactly $6.022\,140\,76 \times 10^{23}$ elementary entities. This number is the fixed numerical value of the Avogadro constant, N_A , when expressed in the unit mol^{-1} and is called the Avogadro number. The amount of substance, symbol n , of a system is a measure of the number of specified elementary entities. An elementary entity may be an atom, a molecule, an ion, an electron, any other particle or specified group of particles.

luminous intensity: *candela* (cd)

It is defined by taking the fixed numerical value of the luminous efficacy of monochromatic radiation of frequency 540×10^{12} Hz, K_{cd} , to be 683 when expressed in the unit $\text{lm} \cdot \text{W}^{-1}$, which is equal to $\text{cd} \cdot \text{sr} \cdot \text{W}^{-1}$, or $\text{cd} \cdot \text{sr} \cdot \text{kg}^{-1} \cdot \text{m}^{-2} \cdot \text{s}^3$, where the kilogram, metre and second are defined in terms of h , c and $\Delta\nu_{\text{Cs}}$.

And then a number of other quantities are defined with these units (most of which we'll be using in coming chapters):

plane angle: *radian* (rad, m/m, the ratio of circular arc length to radius)

solid angle: *steradian* (sr, m^2/m^2 , the ratio of spherical area to square radius)

frequency: *hertz* (Hz, s^{-1})

force: *newton* (N, $\text{kg} \cdot \text{m} \cdot \text{s}^{-2}$)

pressure, stress: *pascal* (Pa, $\text{N}/\text{m}^2 = \text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$)

energy, work, amount of heat: *joule* (J, $\text{N} \cdot \text{m} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$)

power, radiant flux: *watt* (W, $\text{J}/\text{s} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3}$)

electric charge: *coulomb* (C, $\text{A} \cdot \text{s}$)

electric potential difference: *volt* (V, $\text{W}/\text{A} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3} \cdot \text{A}^{-1}$)

capacitance: *farad* (F, $\text{C}/\text{V} = \text{kg}^{-1} \cdot \text{m}^{-2} \cdot \text{s}^4 \cdot \text{A}^2$)

electric resistance: *ohm* (Ω , $\text{V}/\text{A} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3} \cdot \text{A}^{-2}$)

electric conductance: *siemens* (S, $\text{A}/\text{V} = \text{kg}^{-1} \cdot \text{m}^{-2} \cdot \text{s}^3 \cdot \text{A}^2$)

magnetic flux: *weber* (Wb, $\text{V} \cdot \text{s} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} \cdot \text{A}^{-1}$)

magnetic flux density: *tesla* (T, $\text{Wb}/\text{m}^2 = \text{kg} \cdot \text{s}^{-2} \cdot \text{A}^{-1}$)

inductance: *henry* (H, $\text{Wb}/\text{A} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} \cdot \text{A}^{-2}$)

Celsius temperature: *degree Celsius* ($^{\circ}\text{C}$, $\text{K} = ^{\circ}\text{C} + 273.15$)

luminous flux: *lumen* (lm, $\text{cd} \cdot \text{sr} = \text{cd}$)

illuminance: *lux* (lx, $\text{lm}/\text{m}^2 = \text{cd} \cdot \text{m}^{-2}$)

activity referred to a radionuclide: *becquerel* (Bq, s^{-1})

absorbed dose, kerma: *gray* (Gy, $\text{J}/\text{kg} = \text{m}^2 \cdot \text{s}^{-2}$)

dose equivalent: *sievert* (Sv, $\text{J}/\text{kg} = \text{m}^2 \cdot \text{s}^{-2}$)

catalytic activity: *katal* (kat, $\text{mol} \cdot \text{s}^{-1}$)

It is important to pay attention to the units in these definitions. Many errors in calculations can be caught by making sure that the final units are correct, and it can be possible to make a rough estimate of an answer to a problem simply by collecting relevant terms with the right units (this is the subject of *dimensional analysis*). Electromagnetic units are particularly confusing; we will consider them in more detail in Chapter 6. The SI system is also called *MKS* because it bases its units on the meter, the kilogram, and the second. For some problems it will be more convenient to use *CGS* units (based on the

Table 2.2. Selected conversion factors.

1 dyne ($\text{gm} \cdot \text{cm} \cdot \text{s}^{-2}$)	=	1×10^{-5} N
1 erg ($\text{gm} \cdot \text{cm}^2 \cdot \text{s}^{-2}$)	=	1×10^{-7} J
1 horsepower (hp)	=	745.7 W
1 atmosphere (atm)	=	101325 Pa
1 ton (short)	=	2000 pounds
	=	907.18474 kg
1 electron volt (eV)	=	$1.602176462 \times 10^{-19}$ J
1 amu	=	$1.66053873 \times 10^{-27}$ kg
1 angstrom (\AA)	=	1×10^{-10} m
1 fermi (fm)	=	1×10^{-15} m
1 parsec (pc)	=	3.085678×10^{16} m
1 mile (mi)	=	1609.344 m
1 foot (ft)	=	0.3048 m
1 inch (in)	=	0.0254 m
1 liter (L)	=	0.001 m^3
1 pound (lb)	=	.45359237 kg
1 pound-force (lbf)	=	4.44822 N

centimeter, the gram, and the second); MKS is more common in engineering and CGS in physics. A number of other units have been defined by characteristic features or by historical practice; some that will be useful later are given in Table 2.2.

It's often more relevant to know the value on one quantity relative to another one, rather than the value itself. The ratio of two values X_1 and X_2 , measured in *decibels* (dB), is defined to be

$$dB = 20 \log_{10} \frac{X_1}{X_2} \quad . \quad (2.1)$$

If the *power* (energy per time) in two signals is P_1 and P_2 , then

$$dB = 10 \log_{10} \frac{P_1}{P_2} \quad . \quad (2.2)$$

This is because the power is the mean square amplitude (Chapter 3), and so to be consistent with equation (2.1) a factor of 2 is brought in to account for the exponent. An increase of 10 db therefore represents a increase by a factor of 10 in the relative power of two signals, or a factor of 3.2 in their values. A change of 3 dB in power is a change by a factor of 2.

The name decibel comes from Bell Labs. Engineers there working on the telephone system found it convenient to measure the gain or loss of devices on a logarithmic scale. Because the log of a product of two numbers is equal to the sum of their logs, this let them find the overall gain of a system by adding the logs of the components, and using logarithms also made it more convenient to express large numbers. They called the base-10 logarithm the *bel* in honor of Alexander Graham Bell; multiplying by 10 to bring up one more significant digit gave them a tenth-bel, or a decibel.

Some decibel reference levels occur so commonly that they are given names; popular ones include:

- dBV measures an electrical signal relative to 1 volt

- *dBm* measures relative to a 1 mW signal. The power will depend on the (usually unspecified) load, which traditionally is $50\ \Omega$ for radiofrequency signals and $600\ \Omega$ for audio ones (loads will be covered in Chapter 7). In audio recording, this is also called the *Volume Unit* or *VU*.
- *dBspl*, for *Sound Pressure Level* (or just *SPL*), measures sound pressure relative to a reference of 2×10^{-5} Pa, the softest sound that the ear can perceive. The sound of a jet taking off is about 140 dBspl.

Finally, we'll be using a number of fundamental observed constants in nature, including the *gravitational constant* ($G = 6.673(10) \times 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2}$), the *electron mass* ($m_e = 9.10938188(72) \times 10^{-31} \text{ kg}$), and the *proton mass* ($m_p = 1.67262158(13) \times 10^{-27} \text{ kg}$). In this list the digits in parentheses are the standard deviation uncertainty (see Chapter 3) in the corresponding digits, so that for example the error in the value for G is 0.010×10^{-11} (which, compared to the other constants, is an embarrassingly large uncertainty [Gundlach *et al.*, 1996]). Calculating these from first principles remains an open Nobel-prize-worthy problem.

Each fundamental constant can appear in many different types of measurements, and these are done by many different groups, leading to multiple values that unfortunately don't always agree to within their careful error estimates. For this reason, the International Council of Scientific Unions in 1966 formed the *Committee on Data for Science and Technology (CODATA)* to do global optimizations over all these data to come up with an internally-consistent set of values [Mohr *et al.*, 2016].

2.2 PARTICLES AND FORCES

The world is built out of elementary particles and their interactions. There are a number of natural divisions in organization, energy, and length that occur between the structure of the nucleus of an atom and the structure of the universe; it will be useful to briefly survey this range in order to understand the relevant regimes for present and prospective information technologies.

This story starts with quantum mechanics, the laws that govern things that are very small. Around 1900 Max Planck was led by his inability to explain the spectrum of light from a hot oven to propose that the energy of light is quantized in units of $E = h\nu = hc/\lambda$, where ν is the frequency and λ is the wavelength; $h = 6.626 \dots \times 10^{-34} \text{ J} \cdot \text{s}$ is now called *Planck's constant*. From there, in 1905 Einstein introduced the notion of massless photons as the discrete constituents of light, and in 1924 de Broglie suggested that the wavelength relationship applies to massive as well as massless particles by $\lambda = h/p$; λ is the *de Broglie wavelength*, and is a consequence of the *wave-particle duality*: all quantum particles behave as both waves and particles. An electron, or a photon, can diffract like a wave from a periodic grating, but a detector will register the arrival of individual particles. Quantum effects usually become significant when the de Broglie wavelength becomes comparable to the size of an object.

Quantum mechanical particles can be either *fermions* (such as an electron) or *bosons* (such as a photon). Fermions and bosons are as unlike as anything can be in our universe. We will later see that bosons are particles that exist in states that are symmetric under the

interchange of particles, they have an integer spin quantum number, and multiple bosons can be in the same quantum state. Fermions have half-integer spin, exist in states that are antisymmetric under particle interchange, and only one fermion can be in a particular quantum state. Spin is an abstract property of a quantum particle, but it behaves just like an angular momentum (as if the particle is spinning).

Particles can interact through four possible forces: *gravitational*, *electromagnetic*, *weak*, and *strong*. The first two are familiar because they have infinite range; the latter two operate on short ranges and are associated with nuclear and subnuclear processes (the characteristic lengths are approximately 10^{-15} m for the strong force and 10^{-18} m for the weak force). The electromagnetic force is so significant because of its strength: if a quantum atom was held together by gravitational forces alone (like a miniature solar system) its size would be on the order of 10^{23} m instead of 10^{-10} m. The macroscopic forces that we feel, such as the hardness of a wall, are transmitted to us by the electromagnetic force through the electrons in our atoms interacting with electrons in the adjoining atoms in the surface, but can be much more simply described in terms of fictitious effective forces (“the wall is hard”).

All forces were originally thought to be transmitted by an intervening medium, the long-sought *ether* for electromagnetic forces. We now understand that forces operate by the exchange of spin-1 gauge bosons – the photon for the electromagnetic interaction (electric and magnetic fields), the W^\pm and Z^0 bosons for the weak interaction, and eight gluons for the strong interaction (there is not yet a successful quantum theory of gravity). Particle masses arise from interactions with the Higgs field [Aad *et al.*, 2012]. *Quantum ElectroDynamics (QED)* is the theory of the quantum electromagnetic interaction, and *Quantum ChromoDynamics (QCD)* the theory of the strong interaction. The weak and electromagnetic interactions are united in the *electroweak theory*, which, along with QCD is the basis for the *Standard Model*, the current summary of our understanding of particle physics. This amalgam of experimental observations and theoretical inferences successfully predicts most observed behavior extremely accurately, with two important catches: the theory has 20 or so adjustable parameters that must be determined from experiments, and it cannot explain gravitation.

The most fundamental massive particles that we are aware of are the *quarks* and *leptons*. There’s no reason to assume that there’s nothing below them (i.e., turtles all the way down); there’s just not a compelling reason right now to believe that there is. Quarks and leptons appear in the scattering experiments used to study particle physics to be point-particles without internal structure, and are spin-1/2 fermions. The leptons interact through the electromagnetic and weak interactions, and come in pairs: the *electron* and the *electron neutrino* (e^- , ν_e), the *muon* and its neutrino (μ^- , ν_μ), and the *tau lepton* and its neutrino (τ^- , ν_τ). Muons and tau leptons are unstable, and therefore are seen only in accelerators, particle decay products, and cosmic rays. Because neutrinos interact only through the weak force they can pass unhindered through a light-year of lead. They were originally thought to be massless, but their small observed mass is a cosmologically-significant deviation from the Standard Model [Aker *et al.*, 2019].

Quarks interact through the strong as well as weak and electromagnetic interactions, and they come in pairs: *up* and *down*, *charm* and *strange*, and *top* and *bottom*. These fanciful names are just labels for the underlying abstract states. The first member of each pair has charge $+2/3$, the second member has charge $-1/3$, and each charge flavor

comes in three colors (once again, flavor and color are just descriptive names for quantum numbers).

Quarks combine to form *hadrons*; the best-known of which are the two *nucleons*. A proton comprises two ups and a down, and the neutron an up and two downs. The nucleons, along with their excited states, are called *baryons* and are fermions. Transitions between baryon states can absorb or emit spin-1 boson hadrons, called *mesons*. The size of hadrons is on the order of 10^{-15} m, and the energy difference between excited states is on the order of 10^9 electron volts (1 GeV).

The nucleus of an atom is made up of some number of protons and neutrons, bound into ground and excited states by the strong interaction. Typical nuclear sizes are on the order of 10^{-14} m, and energies for nuclear excitations are on the order of 10^6 eV (1 MeV). Atoms consist of a nucleus and electrons bound by the electromagnetic interaction; typical sizes are on the order of 1 angstrom (\AA , 10^{-10} m) and the energy difference between states is on the order of 1 eV. Notice the large difference in size between the atom and the nucleus: atoms are mostly empty space. Atoms can exist in different *isotopes* that have the same number of protons but differing numbers of neutrons, and *ions* are atoms that have had electrons removed or added.

Atoms can bond to form molecules; bond energies are on the order of 1 eV and bond lengths are on the order of 1 \AA . Molecular sizes range from simple diatomic molecules up to enormous biological molecules with 10^6 – 10^9 atoms. Large molecules fold into complex shapes; this progresses from the *primary structure* of a coding sequence, to the *secondary structure* of geometrical motifs, to the *tertiary structure* of functional units, to the *quaternary structure* of molecular assemblies [Goodsell, 2009]. Predicting this folding was one of the most difficult challenges in molecular biology [R.Evans *et al.*, 2018], but has become tractable through machine learning [Jumper *et al.*, 2021].

Macroscopic materials are described by the arrangement of their constituent atoms, and include crystals (which have complete long-range ordering), liquids and glasses (which have short-range order but little long-range order), and gases (which have little short-range order). There are also very interesting intermediate cases, such as quasiperiodic alloys called *quasicrystals* that have deterministic translational order without translational periodicity [DiVincenzo & Steinhardt, 1991], and *liquid crystals* that maintain orientational but not translational ordering [Chandrasekhar, 1992]. Most solids do not contain just a single phase; there are usually defects and boundaries between different kinds of domains.

The atomic weight of an element is equal to the number of grams equal to one mole ($N_A \approx 10^{23}$) of atoms. It is approximately equal to the number of protons and neutrons in an atom, but differs because of the mix of naturally occurring isotopes. 22.4 liters of an ideal gas at a pressure of 1 atmosphere and at room temperature will also contain a mole of atoms. The structure of a material at more fundamental levels will be invisible and can be ignored unless energies are larger than its characteristic excitations. Although we will rarely need to descend below atomic structure, there are a number of important applications of nuclear transitions, such as nuclear power and the use of nuclear probes to characterize materials.

2.3 ORDERS OF MAGNITUDE

Understanding what is possible and what is preposterous requires being familiar with the range of meaningful numbers for each unit; the following lists include some significant ones:

Time

- 10^{-43} s: the Planck time (Problem 2.7)
- 10^{-15} s: this is the period of visible light, and a typical time scale for chemical reactions
- 10^{-12} s: shortest logic gate delay
- 10^{-9} s: atomic excitations and molecular rotations typically have lifetimes on the order of nanoseconds, and this is a characteristic computer clock cycle
- 10^{-3} s: the shortest time difference that is consciously perceptible by people
- 10^{17} s: the approximate age of the observable universe

Power and Energy

- 1 eV: atomic excitations
- 10^6 eV: nuclear excitations
- 10^9 eV: subnuclear excitations
- 10^{28} eV: the Planck energy
- 10 W: laptop computer
- 100 W: workstation; human
- 1000 W: house
- 10^4 W: car
- 10^5 W: building
- 10^7 W: supercomputer
- 10^{26} W: luminosity of the sun
- 10^{-12} W/m²: softest sound that can be heard
- 1 W/m²: loudest sound that can be tolerated
- 10^7 J/kg: energy density of food
- 10^9 J: energy in a ton of TNT
- 10^{20} J: energy consumption in the US per year

Temperature

- 10^{-9} K: laser-cooled gas
- 2.75 K: microwave background radiation from the Big Bang
- 77 K: liquid nitrogen
- 300 K: room temperature
- 6000 K: surface of the sun

Mass

- 10^{-27} kg: proton mass
- 10^{-12} kg: typical cell
- 10^{-5} kg: small insect

10^{16} kg: Earth's biomass
 5.98×10^{24} kg: the mass of the Earth
 10^{42} kg: approximate mass of the Milky Way

Length

10^{-35} m: the Planck distance
 10^{-15} m: size of a proton
 10^{-10} m: size of an atom
 4×10^5 m: height of a Low Earth Orbit satellite above the surface
 6.378×10^6 m: radius of the Earth
 4×10^7 m: height of a geosynchronous satellite above the equator
 10^{11} m: distance from the Earth to the Sun
 10^{20} m: Milky Way radius
 10^{26} m: size of the observable universe

Electromagnetic spectrum

< 0.1 Å: gamma rays
 0.1 – 100 Å: X-rays
 100 – 4000 Å: UV (atomic ionization energy)
 4000 – 7000 Å: visible (this coincides with a transmission band through the atmosphere, and corresponds to 10^{14} – 10^{15} Hz)
 0.7 – 100 μm: IR (thermal radiation)
 0.01 – 10 cm: microwave (GHz)
 0.1 – 10^3 m: radio (MHz–kHz)

Magnetic and Electric Fields

10^{-12} tesla: magnetic field needed for radio reception
 10^{-6} tesla: magnetic field generated by a cordless phone
 3×10^{-5} tesla: magnetic field at the Earth's surface
 20 tesla: large superconducting/normal hybrid magnet
 10^4 A: lightning bolt current
 10^8 V: potential across a lightning bolt
 3×10^6 V/m: breakdown voltage in air

Number

10^5 : number of DNA bases in a bacteriophage
 4×10^6 : bytes in the Bible
 10^9 : number of DNA bases in a mammal
 10^{13} : number of synapses in the human cortex
 10^{21} : bytes passing through the Internet in 2016
 10^{80} : approximate number of atoms in the universe

2.4 SELECTED REFERENCES

[Cohen *et al.*, 2003] Cohen, E. Richard, Lide, David R., & Trigg, George L. (eds). (2003). *A Physicist's Desk Reference*. New York: Springer.

This is a handy summary of units, conversion factors, and governing equations.

[Lerner & Trigg, 2005] Lerner, Rita G., & Trigg, George L. (eds). (2005). *Encyclopedia of Physics*. 3rd edn. New York: Wiley.

A Who's Who of interesting physical phenomena.

[Morrison & Morrison, 1982] Morrison, Philip, & Morrison, Phylis. (1982). *Powers Of Ten: A Book About the Relative Size of Things*. Redding: Scientific American Library.

The Morrisons provide a marvelous tour through the characteristic phenomena at many length scales.

[Nachtmann, 1990] Nachtmann, Otto. (1990). *Elementary Particle Physics: Concepts and Phenomena*. New York: Springer-Verlag. Translated by A. Lahee and W. Wetzel.

A nice introduction to particles and forces.

2.5 PROBLEMS

- (2.1) (a) How many atoms are there in a yoctomole?
 (b) How many seconds are there in a nanocentury? Is the value near that of any important constants?
- (2.2) A large data storage system holds on the order of an exabyte. How tall would a 1 exabyte stack of DVDs be? How does that compare to the distance from Earth to space?
- (2.3) If all the atoms in our universe were used to write an enormous binary number, using one atom per bit, what would that number be (converted to base 10)?
- (2.4) Compare the gravitational acceleration due to the mass of the Earth at its surface to that produced by a 1 kg mass at a distance of 1 m. Express their ratio in decibels.
- (2.5) (a) Approximately estimate the chemical energy in a ton of TNT. You can assume that nitrogen is the primary component; think about what kind of energy is released in a chemical reaction, where it is stored, and how much there is.
 (b) Estimate how much uranium would be needed to make a nuclear explosion equal to the energy in a chemical explosion in 10 000 tons of TNT (once again, think about where the energy is stored).
 (c) Compare this to the *rest mass energy* $E = mc^2$ of that amount of material (Chapter 14), which gives the maximum amount of energy that could be liberated from it.
- (2.6) (a) What is the approximate de Broglie wavelength of a thrown baseball?
 (b) Of a molecule of nitrogen gas at room temperature and pressure? (This requires either the result of Section 3.4.2, or dimensional analysis.)
 (c) What is the typical distance between the molecules in this gas?

-
- (d) If the volume of the gas is kept constant as it is cooled, at what temperature does the wavelength become comparable to the distance between the molecules?
- (2.7) (a) The potential energy of a mass m a distance r from a mass M is $-GMm/r$. What is the *escape velocity* required to climb out of that potential?
- (b) Since nothing can travel faster than the speed of light (Chapter 14), what is the radius within which nothing can escape from the mass?
- (c) If the rest energy of a mass M is converted into a photon, what is its wavelength?
- (d) For what mass does its equivalent wavelength equal the size within which light cannot escape?
- (e) What is the corresponding size?
- (f) What is the energy?
- (g) What is the period?
- (2.8) Consider a pyramid of height H and a square base of side length L . A sphere is placed so that its center is at the center of the square at the base of the pyramid, and so that it is tangent to all of the edges of the pyramid (intersecting each edge at just one point).
- (a) How high is the pyramid in terms of L ?
- (b) What is the volume of the space common to the sphere and the pyramid?
- (This question comes from an entrance examination for humanities students at Tokyo University [*Economist*, 1993].)